

Dedicated to Professor Rudolf Zahradnik on the occasion of his 80th birthday

About non-existence of the molecular ion H_3^{++}

H. Medel-Cobaxin*

*Instituto de Ciencias Nucleares, UNAM,
Apartado Postal 70-543, 04510 México*

A. Alijah†

*Departamento de Química,
Universidade de Coimbra,
3004-535 Coimbra, Portugal*

A.V. Turbiner‡

*Instituto de Ciencias Nucleares, UNAM,
Apartado Postal 70-543, 04510 México*

Abstract

It is shown that the molecular ion H_3^{++} does not exist in a form of the equilateral triangle. To this end, a compact variational method is presented which is based on a linear superposition of six specially tailored trial functions containing non-linear parameters. Careful optimization of a total of fifteen parameters gives consistently lower variational results for the electronic energy than can be obtained with standard methods of quantum chemistry even with huge basis sets as large as mcc-pV7Z.

*Electronic address: medel@nucleares.unam.mx

†Electronic address: alijah@ci.uc.pt

‡Electronic address: turbiner@nucleares.unam.mx

I. INTRODUCTION

The trihydrogen dication, H_3^{++} , which consists of three protons and one electron, is among the simplest Coulomb systems. Its stability has been studied intensely in the sixties and early seventies. In a series of articles, Conroy [1, 2, 3] investigated the potential energy surfaces of the electronic ground state and the lowest excited states at linear and isosceles triangular configurations. He employed a variational approach in which the electronic trial wavefunction is expanded around the center of the nuclear charges. Analyzing the contour plots Conroy concluded that H_3^{++} is not stable. Schwartz and Schaad [4], and Somorjai and Yue [5], who reported single-point calculations of the system (*pppe*) at the supposed equilibrium equilateral triangular configuration of H_3^+ , did not address the stability problem. To assess Conroy's results, Berkowitz and Stocker [6] searched for this ion through charge stripping experiments on H_3^+ . They could not find evidence of stable H_3^{++} . Later, the issue was reconsidered also from the theoretical side, by Shoucri and Darling [7], who examined equilateral configurations with the variational linear combination of atomic orbitals (LCAO) method, and by Hernández and Carbó [8], who studied two particular configurations with a more compact variational approach and obtained total energy values below those published before. No bound state has been determined in these calculations. Johnson and Poshusta [9] reported another single-point calculation in the context of Gaussian basis set optimization for some one-electron systems.

About twenty years later Ackermann *et al.* [10] revisited the question about the existence of H_3^{++} using the finite element method which provided much higher accuracy than previously achieved. The problem of the stability of H_3^{++} was treated keeping the nuclear charge as a continuous parameter. Critical values of the charges for the existence of stable or metastable equilateral triangular configurations were obtained as $Z_c^+ = 0.82$ and $Z_c^+ = 0.95$, respectively. The authors excluded the possibility of stable H_3^{++} in the electronic ground state. However, the explicit electronic energy data are reported only for one particular equilateral triangular configuration at the triangle size $R = 1.68 \text{ a.u.}$. In conclusion, accurate *ab initio* results on the basis of which the non-existence of H_3^{++} can be demonstrated are scarce and not that convincing. This question is thus addressed once again in the present study. One of the motivations of our study is related to a fact that H_3^{++} in equilateral triangular configuration may exist as metastable state in a magnetic field $B \gtrsim 10^8 \text{ G}$ [11].

II. METHODS

We study a Coulomb system of one electron and three protons (*pppe*) which form an equilateral triangle of size R . The protons are assumed to be infinitely massive according to the Born-Oppenheimer approximation at zero order. The Schrödinger equation for the system is written as

$$\left[\mathbf{p}^2 + \frac{6}{R} - \frac{2}{r_1} - \frac{2}{r_2} - \frac{2}{r_3} \right] \Psi(\mathbf{r}) = E\Psi(\mathbf{r}) , \quad (1)$$

where $\mathbf{p} = -i\nabla$ is the electron momentum, r_1, r_2 and r_3 are the distances from each proton to the electron and R is the interproton distance, see Figure 1. Atomic units are used throughout ($\hbar=m_e=e=1$), although energies are expressed in Rydbergs (Ry).




FIG. 1: Molecular ion H_3^{++} in equilateral triangular configuration. Protons are fixed on the vertexes (on the x-y plane). Point **e** represents the electron position.

Our goal is to study the stability of the molecular ion H_3^{++} . If such an ion exists, it implies the existence of the ground state of the system (*pppe*). Based on symmetry arguments it seems evident that the optimal geometry of (*pppe*) in the case of existence of a bound state is the equilateral triangle.

Two methods are used to explore the system: (i) variational with physically relevant trial functions (see e.g. [11]) which we will call *specialized* and (ii) *standard* variational based on using standard Gaussian trial functions as implemented in *ab initio* quantum chemistry packages such as MOLPRO [12]. Both methods lead to highly accurate quantitative results for total energy versus the size of the triangle.

In the first variational approach, a trial function is taken in a form of linear superposition

of six basis functions

$$\Psi_{trial} = \sum_{j=1}^6 A_j \psi^{(j)} , \quad (2)$$

where A_j are linear parameters. Each function $\psi^{(j)}$ is chosen in such a way as to describe different physical characteristics of the system. In general, $\psi^{(j)}$ has the form of a symmetrized product of three Coulomb orbitals

$$\psi_C = e^{-\alpha_1 r_1 - \alpha_2 r_2 - \alpha_3 r_3} . \quad (3)$$

Let us give a brief description of each of them:

$\psi^{(1)}$: All α 's are chosen to be equal to α_1 ,

$$\psi^{(1)} = e^{-\alpha_1 (r_1 + r_2 + r_3)} . \quad (4)$$

It is a Heitler-London type function. This corresponds to *coherent* interaction between the electron and all protons. Supposedly, it describes the system at small interproton distances and, probably, the equilibrium configuration. It might be verified a posteriori.

$\psi^{(2)}$: Two α 's are equal to zero and the remaining one is set to be equal to α_2 ,

$$\psi^{(2)} = e^{-\alpha_2 r_1} + e^{-\alpha_2 r_2} + e^{-\alpha_2 r_3} . \quad (5)$$

It is a Hund-Mulliken type function. This function possibly describes the system at large distances, where essentially the electron interacts with only one proton at a time thus realizing *incoherent* interaction.

$\psi^{(3)}$: One α is equal to zero, two others are different from zero and equal to each other and to α_3 ,

$$\psi^{(3)} = e^{-\alpha_3 (r_1 + r_2)} + e^{-\alpha_3 (r_1 + r_3)} + e^{-\alpha_3 (r_2 + r_3)} . \quad (6)$$

It is assumed that this function describes the system H_2^+ plus proton when a triangle is of a sufficiently small size. In fact, it is the Heitler-London function of H_2^+ symmetrized over protons.

$\psi^{(4)}$: One α is equal to zero and two others are different from each other being equal to $\alpha_{4,5}$, respectively,

$$\begin{aligned}\psi^{(4)} = & e^{-\alpha_4 r_1 - \alpha_5 r_2} + e^{-\alpha_4 r_2 - \alpha_5 r_1} + e^{-\alpha_4 r_1 - \alpha_5 r_3} \\ & + e^{-\alpha_4 r_3 - \alpha_5 r_1} + e^{-\alpha_4 r_2 - \alpha_5 r_3} + e^{-\alpha_4 r_3 - \alpha_5 r_2} .\end{aligned}\quad (7)$$

It is assumed that this function describes the system H_2^+ plus one proton. In fact, it is the Guillemin-Zener function of H_2^+ symmetrized over protons. If $\alpha_4 = \alpha_5$, the function $\psi^{(4)}$ is reduced to $\psi^{(3)}$. If $\alpha_4 = 0$, the function $\psi^{(4)}$ is reduced to $\psi^{(2)}$. Hence $\psi^{(4)}$ is a non-linear interpolation between $\psi^{(2)}$ and $\psi^{(3)}$. It has to describe intermediate interproton distances.

$\psi^{(5)}$: Two α 's are equal but the third one is different,

$$\begin{aligned}\psi^{(5)} = & e^{-\alpha_6 r_1 - \alpha_6 r_2 - \alpha_7 r_3} + e^{-\alpha_6 r_1 - \alpha_6 r_3 - \alpha_7 r_2} \\ & + e^{-\alpha_6 r_2 - \alpha_6 r_1 - \alpha_7 r_3} + e^{-\alpha_6 r_2 - \alpha_6 r_3 - \alpha_7 r_1} \\ & + e^{-\alpha_6 r_3 - \alpha_6 r_1 - \alpha_7 r_2} + e^{-\alpha_6 r_3 - \alpha_6 r_2 - \alpha_7 r_1} .\end{aligned}\quad (8)$$

It describes a “mixed” state of three hydrogen atoms. If $\alpha_6 = \alpha_7$, the function $\psi^{(5)}$ is reduced to $\psi^{(1)}$. If $\alpha_6 = 0$, the function $\psi^{(5)}$ is reduced to $\psi^{(2)}$. If $\alpha_7 = 0$, the function $\psi^{(5)}$ is reduced to $\psi^{(3)}$. Hence $\psi^{(5)}$ is a non-linear interpolation between $\psi^{(1)}$, $\psi^{(2)}$ and $\psi^{(3)}$. As function (7) this is a type of Guillemin-Zener function and should describe intermediate interproton distances.

$\psi^{(6)}$: All α 's are different,

$$\begin{aligned}\psi^{(6)} = & e^{-\alpha_8 r_1 - \alpha_9 r_2 - \alpha_{10} r_3} + e^{-\alpha_8 r_1 - \alpha_9 r_3 - \alpha_{10} r_2} \\ & + e^{-\alpha_8 r_2 - \alpha_9 r_1 - \alpha_{10} r_3} + e^{-\alpha_8 r_2 - \alpha_9 r_3 - \alpha_{10} r_1} \\ & + e^{-\alpha_8 r_3 - \alpha_9 r_1 - \alpha_{10} r_2} + e^{-\alpha_8 r_3 - \alpha_9 r_2 - \alpha_{10} r_1} .\end{aligned}\quad (9)$$

This is a general non-linear interpolation of all functions $\psi^{(1-5)}$.

The total number of the parameters of the function (2) is equal to 15, where five of them are linear ones. Note that A_6 can be fixed at $A_6 = 1$.

In standard *ab initio* calculations, Ψ_{trial} is most commonly expanded in terms of Gaussian basis functions [13] $\psi_{I,i}$ centered at atoms $I = 1, 2, 3$,

$$\Psi_{trial} = \sum_{I=1}^3 \sum_{i=1}^n \psi_{I,i}(\mathbf{r}_I) c_{Ii} \quad (10)$$

whose coefficients c_{Ii} are then determined variationally. The basis functions ψ_{Ii} themselves are built up by primitive Gaussians [14]

$$\psi_{I,i=\{n,l,m\}}(\mathbf{r}_I) = \sum_{\kappa} N Y_{lm}(\theta, \phi) r^{2n-2-l} e^{-\alpha_{\kappa} r^2} d_{\kappa} \quad (11)$$

with contraction coefficients d_{κ} held fixed.

Our calculations were performed using the Hartree-Fock code implemented in the MOLPRO suite of programs [12] with the correlation consistent cc-pV6Z and modified mcc-pV7Z basis sets [15, 16]. The cc-pV6Z basis set contains 91 contracted Gaussians per atom, with ℓ quantum numbers up to $\ell = 5$, i.e. [6s 5p 4d 3f 2g 1h], yielding a total of 273 basis functions. The mcc-pV7Z basis includes $\ell = 6$ functions, leading to 140 contracted Gaussians per atom, [7s 6p 5d 4f 3g 2h 1i], or 420 basis functions in total. Calculations were carried out for a range of equilateral triangular configurations using C_s symmetry. In this point group, the lowest electronic state is $^2A'$. The total number of contracted Gaussians of this symmetry is 168 for the cc-pV6Z basis set and 255 for the mcc-pV7Z basis set, respectively. The cc-pV6Z results, which are not reported here explicitly, have been generated to assess the accuracy of this type of calculations. Judging from such a comparison, we estimate the accuracy of the mcc-pV7Z data to about $10^{-5} a.u.$ over a large range of distances, deteriorating somewhat at short distances where the basis functions tend to become linearly dependent.

III. RESULTS

In framework of the specialized variational method (i) some numerical computations were made. The minimization routine MINUIT [17] from the CERN-LIB library was used as well as D01FCF routine from the NAG-LIB [18] for three-dimensional numerical integration. Numerical values of the total energy E_T of the system (*pppe*) for different values of the interproton distance R were obtained, see Table I. The results from the MOLPRO calculation with a huge standard-type basis set (mcc-pV7Z) are given for comparison. A problem of the standard approach is its slow convergence with respect to the angular momentum quantum

number ℓ , requiring the use of large basis sets. The method based on the specially tailored trial function, Eq. (2), leads to systematically lower variational energy values with considerably less terms. It should be noted that this method relies on a careful optimization of non-linear parameters.

Size $R(\text{a.u.})$	Total Energy $E_T(\text{Ry})$	
	Variational, specialized	Variational, standard
0.10	51.2605684	51.2608280
0.20	21.7652245	21.7658978
0.30	12.3060338	12.3066419
0.40	7.8158809	7.8161665
0.50	5.2770999	5.2772488
0.60	3.6886676	3.6887578
0.70	2.6262531	2.6263119
0.80	1.8810732	1.8811131
0.90	1.3394273	1.3394556
1.00	0.9346477	0.9346656
1.10	0.6253608	0.6253729
1.20	0.3847163	0.3847301
1.30	0.1946538	0.1946668
1.40	0.0426368	0.0426474
1.50	-0.0802535	-0.0802489
1.65	-0.2237211	-0.2237116
2.00	-0.4355913	-0.4355854
2.50	-0.5827041	-0.5826899
3.00	-0.6486047	-0.6485824
3.50	-0.6793259	-0.6793070

TABLE I: Variational results obtained with the specialized method (2) and with a standard quantum chemistry method MOLPRO for the total energy E_T as a function of the internuclear distance R for the system *pppe* in the equilateral triangular geometry. For $R = 3.50$, in [8] $E_T = -0.678$.

Different studies have been done for $R = 1.68$ a.u. This distance corresponds to an early

estimate of the equilibrium distance (R_e) for the molecular ion H_3^+ in triangular equilateral configuration [2]. It provides a natural explanation why this was considered. Nowadays it is known with high accuracy that the equilibrium distance for H_3^+ is $R = 1.65$ [19]. We present, in Table II, a comparison of the results obtained at $R = 1.68$ by the standard method of quantum chemistry, LCAO, which uses optimized Gauss-type (GTO) or Slater-type (STO) atomic orbitals, the Finite Element method, the MOLPRO calculation with the massive aug-mcc-pVZ basis set [16] consisting of 567 basis functions, 336 of which have symmetry A' , and the variational approach, Eq. (2).

Author	Method	Energy(Ry)
Conroy [1]	STO	-0.2431714
Johnson and Poshusta [9]	GTO	-0.24748, est. exact -0.248
Schwartz and Schaad [4]	Variational	-0.24746
Somorjai and Yue [5]	Variational	-0.23416
Ackermann <i>et al.</i> [10]	F. E.	-0.2477134
Present work	MOLPRO, aug-mcc-pV7Z	-0.2477064
Present work	Variational (2)	-0.2477132

TABLE II: Comparison different results for E_T at $R = 1.68$ a.u.

Plotting the data results in a smooth monotonous curve of the total Energy E_T as a function of the internuclear distance R , see Fig. 2.

As a conclusion we have to state that the total energy E_T as a function of the internuclear distance R does not indicate either to a minimum or even slight non-adiabatic irregularities for finite R . Thus, the molecular ion H_3^{++} does not exist in equilateral triangular configuration in the Born-Oppenheimer approximation.

Acknowledgments

HMC expresses his deep gratitude to J.C. López Vieyra for the valuable comments and for their interest in the present work, AVT thanks Universite Libre de Bruxelles for the hospitality extended to him where this work was completed. This work was supported in




FIG. 2: E_T as function of the internuclear distance R .

part by FENOMECC and PAPIIT grant **IN121106** (Mexico).

-
- [1] H. Conroy, J. Chem. Phys. 41, 1327 (1964)
 - [2] H. Conroy, J. Chem. Phys. 41, 1341 (1964)
 - [3] H. Conroy, J. Chem. Phys. 51, 3979 (1969)
 - [4] M. E. Schwartz and L. J. Schaad, J. Chem. Phys. 47, 5325 (1967)
 - [5] R. L. Somorjai and C. P. Yue, J. Chem. Phys. 53, 1657 (1970)
 - [6] E. H. Berkowitz and H. Stocker, J. Chem. Phys. 55, 4606 (1971)
 - [7] R. M. Shoucri and B. T. Darling, J. Chem. Phys. 56, 1789 (1972)
 - [8] J. A. Hernández and R. Carbó, J. Chem. Phys. 62, 2637 (1975)
 - [9] J. W. Johnson and R. D. Poshusta, Int. J. Quantum Chem. 11, 885 (1977)
 - [10] J. Ackermann, B. Erdmann and R. Roitzsch, J. Chem. Phys. 101, 7643 (1994)
 - [11] A.V. Turbiner and J.C. Lopez Vieyra, *Phys.Rev.* **A66**, 023409 (2002)

- [12] H.-J. Werner, P. J. Knowles, R. Lindh, F. R. Manby, M. Schütz, P. Celani, T. Korona, G. Rauhut, R. D. Amos, A. Bernhardsson, A. Berning, D. L. Cooper, M. J. O. Deegan, A. J. Dobbyn, F. Eckert, C. Hampel, G. Hetzer, A. W. Lloyd, S. J. McNicholas, W. Meyer, M. E. Mura, A. Nicklass, P. Palmieri, R. Pitzer, U. Schumann, H. Stoll, A. J. Stone, R. Tarroni, and T. Thorsteinsson, MOLPRO, version 2006.1, a package of ab initio programs (2006), see <http://www.molpro.net>
- [13] S. F. Boys, Proc. R. Soc., London **A200**, 542 (1950)
- [14] S. Huzinaga, J. Chem. Phys. **42**, 1293 (1965)
- [15] T. H. Dunning, Jr., J. Chem. Phys. **90**, 1007 (1989)
- [16] S. T. Mielke, B. C. Garrett, and K. A. Peterson, J. Chem. Phys. **116**, 4142 (2002)
- [17] CERN Program Library Long Writeup D506. MINUIT. *Function Minimization and Error Analysis* Reference Manual, Version 94.1, F. James, CERN Geneva, Switzerland (1994-1998)
- [18] NAG Fortran Library Mark 19
- [19] W. Meyer, P. Botschwina and P. Burton, J. Chem. Phys. **84**, 891 (1986)